

Table des matières

1. Systematic Reviews and Meta-Analysis	1
1.1. Generic Acupuncture	1
1.1.1. Lin 2024	1
1.1.2. Hou 2023	2
1.1.3. Huang 2023 (cerebral small vessel disease)	2
1.1.4. Li 2023	3
1.1.5. Wang 2023	4
1.1.6. Yin 2023	4
1.1.7. Su 2021 ☆	5
1.1.8. Jiao 2020	6
1.1.9. Li 2020	6
1.1.10. Min 2019	7
1.1.11. Min 2016 (Vascular Cognitive Impairment) ☆	8
1.1.12. Deng 2016 ☆	8
1.1.13. Lu 2015 (Vascular Cognitive Impairment) ☆☆	9
1.1.14. Hu 2014 ☆☆	9
1.1.15. Zhou 2013 (Vascular Cognitive Impairment) ☆☆	10
1.1.16. Cao 2013 Ø	10
1.1.17. Leung 2013 ~	11
1.2. Special Acupuncture Techniques	11
1.2.1. Acupuncture combined with physical therapies	11
1.2.1.1. Li 2021	11
1.2.2. Comparison of Acupuncture techniques	12
1.2.2.1. Yi 2024	12
1.2.3. Electroacupuncture	13
1.2.3.1. Shen 2024	13
1.2.3.2. Kim 2019 ☆	13
Scalp Acupuncture	14
1.2.4.1. Mai 2015	14
1.2.5. Auricular Acupuncture	15
1.2.5.1. Kwon 2018	15
1.2.6. Acupression	15
1.2.6.1. Liu 2018 ☆	15
1.3. Special Clinical Forms	16
1.3.1. Poststroke Cognitive Impairment	16
1.3.2. Sleep Disturbance in Mild Cognitive Impairment	16
1.3.2.1. O'Caomh 2019	16
2. Overviews of Systematic Reviews	17
2.1. Han 2022	17
2.2. He 2021	17
2.3. Hou 2020 (Methodology and Reporting Quality)	18
3. Clinical Practice Guidelines	19
3.1. Istituto Superiore di Sanità (ISS, Italy) 2024 Ø	19

Cognitive impairment

Troubles cognitifs

1. Systematic Reviews and Meta-Analysis

1.1. Generic Acupuncture

1.1.1. Lin 2024

Lin G, Yie SLJ, Guo S, Li X, Xu L. Clinical evidence of acupuncture for amnesic mild cognitive impairment: A systematic review and meta-analysis of randomized controlled trials. *Complement Ther Med*. 2025 Mar;88:103114. <https://doi.org/10.1016/j.ctim.2024.103114>

Background	People with amnesic mild cognitive impairment (aMCI) carry a substantial risk of developing dementia compared to non-amnesic MCI (naMCI). Several previous studies proved the remarkable effectiveness of acupuncture for MCI, but they didn't distinguish between aMCI and naMCI. We conducted this meta-analysis to systematically assess the evidence of the efficacy of acupuncture in this unique population with aMCI.
Methods	We comprehensively searched nine databases on January 09, 2024, to identify relevant articles estimating the effects of acupuncture for aMCI, and then assessed the risk of bias of the included trials utilizing the RoB 2.0 tool which included the domain of randomization process, deviation from intended interventions, missing outcome data, measurement of the outcome, selection of the reported outcome, and overall bias. The results of this meta-analysis were exhibited with forest plots. Sensitivity analyses were conducted to determine the robustness of the pooled results, and publication bias was estimated by Egger's and Begg's tests. Besides, we also performed subgroup analysis to determine whether there was a difference in therapeutic effects between four weeks and eight weeks of treatment duration. The certainty of the evidence was graded using GRADEpro GDT.
Results	A total of 15 randomized controlled trials (RCTs) involving 908 people with aMCI were included in this study. According to the meta-analysis, acupuncture treatment provided a remarkable improvement in cognitive function as assessed by Mini-Mental State Examination (MD = 1.09, 95 %CI [0.86, 1.31], $p < 0.00001$), Montreal Cognitive Assessment (MD = 0.93, 95 %CI [0.80, 1.07], $p < 0.00001$), Alzheimer's Disease Assessment Scale-Cognitive (MD = 1.00, 95 %CI [-1.23, -0.77], $p < 0.00001$), and P300 latency (MD = -15.40, 95 %CI [-23.68, -7.12], $p = 0.0003$). Subgroup analysis showed evidence that the efficacy of four weeks of acupuncture treatment was consistent with that of eight weeks. Sensitivity analyses, Egger's and Begg's tests suggested the pooled results were robust and reliable. The overall quality of the evidence, as appraised by the GRADE criteria, was very low or low for all outcomes.
Conclusions	The evidence from 15 RCTs demonstrated that acupuncture interventions are effective in ameliorating cognitive function in people with aMCI. There is a need for larger-scale multicenter RCTs using standardised training protocols and more rigorous designs to confirm the conclusions further.

1.1.2. Hou 2023

Hou Z, Yu X, Chen J, Brenner JS, Sun Z, Shang H. Does acupuncture have advantages in the rehabilitation of vascular mild cognitive impairment? A systematic review and meta-analysis. *Heliyon*. 2023 Aug 9;9(8):e18955. <https://doi.org/10.1016/j.heliyon.2023.e18955>

Background	Vascular mild cognitive impairment (VMCI) is a common impairment caused by vascular factors. VMCI often occurs after stroke, and it is the main clinical manifestation of long-term disability. Many patients are treated with acupuncture in combination with other therapies. However, evidence regarding the effectiveness of this treatment regimen is lacking.
Aims	This meta-analysis aimed to evaluate the efficacy of acupuncture therapy for treating VMCI.
Methods	This systematic review was conducted in accordance with the preferred reporting and meta-analysis guidelines. The CNKI, Wanfang, VIP, CBM, Cochrane Library, PubMed and Embase databases were searched from inception to August 20, 2022. After two researchers independently screened the literature, they extracted the data and evaluated the risk of bias in the included studies. Revman 5.3 software was used for the meta-analysis.
Summary of review	Thirty-two randomized controlled trials (RCTs) were included. The overall effective rate of acupuncture for treating VMCI was 3.06, 95% CI [2.39, 3.91], (P < 0.05). Montreal Cognitive Assessment (MoCA), Mini-Mental State Examination (MMSE), Barthel Index and Activities of Daily Living (ADLs) scores significantly differed between the treatment and control groups, with weighted mean differences (WMDs) [95% CI] (P value) of 1.97 [1.44, 2.49] (P < 0.05), 2.02 [1.50, 2.54] (P < 0.05), 5.54 [3.81, 7.28] (P < 0.05), and 3.43 [2.53, 4.33] (P < 0.05), respectively. The overall effective rate of electroacupuncture (EA) for treating VMCI was better than that of the control group (RR = 2.25, 95% CI, [1.13, 4.50], P < 0.05). MoCA, MMSE, Barthel index and ADL scores differed significantly between the treatment and control groups, with WMDs [95% CI] (P value) of 1.79 [1.20, 2.38] (P < 0.05), 1.45 [0.87, 2.03] (P < 0.05), 5.78 [2.38, 9.18] (P < 0.05), and 3.15 [2.15, 4.15] (P < 0.05), respectively. Acupuncture alone and combined with drug therapy were thus superior to drug therapy alone for improving cognitive function. EA also has potential advantages.
Conclusions	Acupuncture combined with another therapy is better than other therapies alone, such as simple drug therapy, for treating VMCI. However, variations in study duration (4-12 weeks) limit us from drawing any definitive conclusions about long-term effects. Therefore, more RCTs with rigorous designs and reasonable treatment and follow-up durations are needed.

1.1.3. Huang 2023 (cerebral small vessel disease)

Huang XT, Chen CY, Zhang QF, Lu LH, She YL, Fang XY. Meta-analysis of the efficacy of acupuncture in the treatment of the vascular cognitive impairment associated with cerebral small vessel disease. *Explore (NY)*. 2023 Jul-Aug;19(4):509-518. <https://doi.org/10.1016/j.explore.2022.10.019>

Objective	To systematically evaluate the efficacy and safety of acupuncture in the treatment of the vascular cognitive impairment (VCI) associated with cerebral small vessel disease (CSVD-VCI) and to provide a theoretical basis for clinical acupuncture treatment for CSVD-VCI.
------------------	--

Method	Various databases, including China National Knowledge Infrastructure, Wanfang Data, Chinese Science and Technology Journal Database, Chinese BioMedical Literature Service System, PubMed, the Cochrane Library, and EBSCOhost, were searched for randomized controlled trials (RCTs) related to acupuncture treatment for CSVD-VCI. The quality of the included trials was evaluated, and a meta-analysis was conducted using the Review Manager 5.4 software.
Results	Ten articles on RCTs were included, involving 761 patients, i.e., 381 in the acupuncture group and 380 in the control group. The meta-analysis results indicated that the use of acupuncture alone and acupuncture alongside other therapies for CSVD-VCI could improve the overall clinical response rate [odds ratio = 3.51, 95% confidence interval (CI) = (2.05, 6.00), $P < 0.00001$], increase the patients' Montreal Cognitive Assessment scores [mean difference (MD) = 3.33, 95%CI (2.98, 3.68), $P < 0.00001$], Mini-Mental State Examination scores [MD = 2.78, 95%CI (2.51, 3.06), $P < 0.00001$], and activities of daily living scores [MD = 6.30, 95%CI (4.22, 8.37), $P < 0.00001$], and shorten the latency of auditory evoked potential P300 [MD = -14.67, 95%CI (-19.54, -9.80), $P < 0.00001$].
Conclusion	Acupuncture alone and acupuncture alongside other therapies are superior to non-acupuncture-based therapies in the treatment of CSVD-VCI. However, due to the small number of relevant available articles and their general low quality, this conclusion may be biased. More clinical RCTs with a larger sample size and higher quality are needed to support this theory.

1.1.4. Li 2023

Li R, Xu C, Zhong P, Wang K, Luo Y, Xiao L, Dai X, Han J, Zhang X. Efficacy of acupuncture and pharmacological therapies for vascular cognitive impairment with no dementia: a network meta-analysis. *Front Aging Neurosci.* 2023 Jun 15;15:1181160. <https://doi.org/10.3389/fnagi.2023.1181160> 37396654; PMID: PMC10310406.

Background and objective	Vascular cognitive impairment with no dementia (VCIND) is considered to be the prodromal stage of vascular dementia, characterized by insidious onset. Although acupuncture and drug therapies are effective, the optimal therapy for VCIND remains to be further determined. Therefore, we conducted a network meta-analysis to compare the effectiveness of acupuncture therapies and current common medicines for VCIND.
Methods	We searched eight electronic databases to identify eligible randomized controlled trials of patients with VCIND treated by acupuncture or drug therapies. The primary outcome was Montreal Cognitive Assessment, and the secondary outcome was Mini-Mental State Examination. We conducted the network meta-analysis within a Bayesian framework. Weighted mean difference with 95% confidence intervals were applied as effect sizes to continuous data for all outcomes. Sensitivity analysis was done to assess the robustness of the findings, and we also carried out a subgroup analysis based on age. We assessed the risk of bias using the Risk of Bias 2.0 tool and applied the Grade of Recommendation Assessment, Development and Evaluation (GRADE) to assess the quality of the outcomes. This study was registered with PROSPERO, number CRD42022331718.

Results	A total of 33 studies with 14 interventions were included, including 2603 participants. In terms of the primary outcome, manual acupuncture plus herbal decoction was considered to be the most effective intervention (P = 91.41%), followed by electroacupuncture (P = 60.77%) and manual acupuncture plus piracetam (P = 42.58%), whereas donepezil hydrochloride ranked the least efficacious intervention (P = 54.19%). For the secondary outcome, electroacupuncture plus nimodipine was considered to be the most effective intervention (P = 42.70%), followed by manual acupuncture plus nimodipine (P = 30.62%) and manual acupuncture (P = 28.89%), whereas nimodipine ranked the least efficacious intervention (P = 44.56%).
Conclusion	Manual acupuncture plus herbal decoction might be the most effective intervention for VCIND. The combination of acupuncture and drug therapy had a tendency to perform better than monotherapy in terms of clinical outcomes.

1.1.5. Wang 2023

Wang YY, Wang XX, Chen L, Liu Y, Li YR. A systematic review and network meta-analysis comparing various non-pharmacological treatments for older people with mild cognitive impairment. *Asian J Psychiatr.* 2023 Aug;86:103635. <https://doi.org/10.1016/j.ajp.2023.103635>

Background and objective	Non-pharmacological therapy appeared to alleviate Mild Cognitive Impairment (MCI) symptoms and signs, according to systematic studies. This network meta-analysis aimed to assess the impact of non-pharmacological therapies on improving cognition in individuals with MCI and identified the most effective intervention.
Methods	We reviewed six databases in search of potentially relevant studies of non-pharmacological therapies such as Physical exercise (PE), Multidisciplinary intervention (MI), Musical therapy (MT), Cognitive training (CT), Cognitive stimulation (CS), Cognitive rehabilitation (CR) Art therapy (AT), general psychotherapy or interpersonal therapy (IPT), and Traditional Chinese Medicine (TCM) (such as acupuncture therapy, massage, auricular-plaster and other related systems) and others. Excluded the literature such as missing full text, missing search results, or no reporting specific values and combined with the inclusion criteria and exclusion criteria in this article, the literature ultimately included in the analysis addressed the following seven non-drug therapies PE, MI, MT, CT, CS, CR, AT. Mini-mental state evaluation paired meta-analyses were undertaken by taking weighted average mean differences with confidence intervals (CI) of 95%. The network meta-analysis was conducted to compare various therapies.
Results	A total of 39 randomized controlled trials, including two three-arm studies, with 3157 participants were included. PE was most likely to be the most effective intervention to slow down the cognitive ability of patients (SMD = 1.34, 95%CI: 0.80, 1.89). CS and CR had no significant effect on cognitive ability.
Conclusions	The non-pharmacological therapy had the potential to greatly promote the cognitive ability of the adult population with MCI. PE had the best chance of being the best non-pharmacological therapy. Due to the limited sample size, substantial variability among different study designs, and the potential for bias, the results should be regarded with caution. Our findings should be confirmed by future multi-center randomized controlled, high-quality large-scale studies.

1.1.6. Yin 2023

Yin Z, Li Y, Jiang C, Xia M, Chen Z, Zhang X, Zhao L, Liang F. Acupuncture for mild cognitive impairment: A systematic review with meta-analysis and trial sequential analysis. *Front Neurol.* 2023 Jan 6;13:1091125. <https://doi.org/10.3389/fneur.2022.1091125>

Background	There is insufficient evidence to support the use of acupuncture for mild cognitive impairment (MCI), and there is no consensus on its efficacy. This review aimed to determine the acupuncture effect in patients with MCI.
Methods	Relevant and potentially eligible randomized controlled trials (RCTs) of acupuncture for MCI were obtained from four Chinese databases, four English databases, and additional resources up to 1 August 2022. The primary outcome was the improvement in overall cognitive function (OCF). Secondary outcomes were improved memory function (MF) and activities of daily living (ADLs). The revised Cochrane collaboration risk of bias (ROB) assessment tool (ROB 2.0) was applied to evaluate their methodological quality. The Review Manager software v 5.4 was used for analyses. Trial sequential analysis (TSA) 0.9.5.10 β software was used to estimate the required sample size and test the reliability of the pooled outcome. The quality of evidence was assessed using the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) tool.
Results	This meta-analysis included 11 RCTs with a total of 602 patients . The methodological quality of all trials was moderate. Low-quality evidence showed that acupuncture significantly improved OCF (Mini-Mental State Examination (MMSE): mean difference (MD) = 1.22, 95% confidence interval (CI): 0.78-1.66; the Montreal Cognitive Assessment Scale (MoCA): MD = 1.22, 95% CI: 0.47-1.97). In subgroup analyses, it was revealed that acupuncture significantly increased OCF in patients with MCI when compared to conventional medicine (CM) and sham acupuncture (SA). TSA's findings indicated that the evidence of improving OCF with acupuncture for patients with MCI was conclusive. Meanwhile, there is no statistical difference in the improvement of MF and ADL between acupuncture and CM. TSA showed that the evidence of improving MF and ADL for patients who had MCI and received acupuncture was inconclusive. The shreds of evidence of improving MF and ADL were ranked from low to critically low.
Conclusion	Acupuncture appears to be an effective clinical application method for improving OCF in patients with MCI. However, due to low-quality evidence, more relevant and high-quality research is needed in this field.

1.1.7. Su 2021 ☆

Su XT, Sun N, Zhang N, Wang LQ, Zou X, Li JL, Yang JW, Shi GX, Liu CZ. Effectiveness and Safety of Acupuncture for Vascular Cognitive Impairment: A Systematic Review and Meta-Analysis. *Front Aging Neurosci.* 2021. [221283]. <https://doi.org/10.3389/fnagi.2021.692508>

Background	Acupuncture may be a promising complementary therapy for vascular cognitive impairment (VCI) and has been extensively applied in China. However, its potential effects remain uncertain, and the clinical findings are inconsistent. This review aimed to systematically appraise the overall effectiveness and safety of acupuncture in treating VCI.
Methods	To investigate the effects of acupuncture on VCI from inception to February 28, 2021 using randomized clinical trials (RCTs), seven electro-databases [Cochrane Central Register of Controlled Trials (CENTRAL), PubMed, Embase, China National Knowledge Infrastructure (CNKI), Chinese Biomedical Literature Database (CBM), VIP, and Wanfang] were searched. Two independent investigators identified the eligible RCTs and extracted data into predesigned forms. The risk of bias (ROB) within each individual trial was evaluated using the Cochrane Collaboration's tool. Meta-analyses were conducted for calculating comparative effects in the RevMan software (version 5.3). The strength of attained evidence was rated using the online GRADEpro approach.

Results	A total of 48 RCTs involving 3,778 patients with VCI were included. The pooled data demonstrated that acupuncture was more beneficial for a global cognitive function [mean difference (MD) 1.86, 95% CI 1.19-2.54, $p < 0.01$] and activities of daily living (MD -3.08, 95% CI -4.81 to -1.35, $p < 0.01$) compared with western medicine (WM). The favorable results were also observed when acupuncture was combined with WM (MD 2.37, 95% CI 1.6-3.14, $p < 0.01$) or usual care (UC, MD 4.4, 95% CI 1.61-7.19, $p = 0.002$) in comparison with the corresponding control conditions. Meanwhile, the subgroup analysis did not indicate a statistical effect difference between manual acupuncture (MA) and electroacupuncture (EA) (inter-group $I^2 < 50\%$ and $p > 0.1$) when comparing acupuncture with WM. There were no significant differences in the occurrence of adverse events (AEs) between the acupuncture group and the control group ($p > 0.05$). Owing to the poor methodological quality and considerable heterogeneity among studies, the certainty of the evidence was low or very low.
Conclusions	This review suggests that acupuncture as a monotherapy or an adjuvant therapy may play a positive role in improving the cognition and daily performance of VCI patients associated with few side effects. The difference in styles may not significantly influence its effectiveness. More rigorously designed and preregistered RCTs are highly desirable to verify the therapeutic benefits and determine an optimal acupuncture paradigm. The methodological and reporting quality of future researches should be enhanced by adhering to authoritative standardized statements.

1.1.8. Jiao 2020

Jiao Lan, Ji-fei Miao, Shu-qi Ge, Tie-qu Chai, ... Li-ming Lu. Acupuncture for cognitive impairment in vascular dementia, Alzheimer's disease and mild cognitive impairment: A systematic review and meta-analysis. *European Journal of Integrative Medicine*. 2020;35. [212068]. [doi](#)

Introduction	Cognitive impairment is a worldwide health problem. Numerous studies have been conducted to evaluate the effect of acupuncture on cognitive impairment. However, it is still unclear that if acupuncture shows the same efficacy on cognitive impairment caused by different diseases. Therefore, we conducted a systematic review and meta-analysis based on the current evidence to evaluate the efficacy and safety of acupuncture for cognitive impairment in vascular dementia (VD), Alzheimer's disease (AD) and mild cognitive impairment (MCI) patients.
Methods	Five databases were searched from their inception to December 2019. Randomized controlled trials (RCTs) involving VD, AD or MCI treated by acupuncture alone or as part of combination therapy were included. The primary outcomes were the Mini-Mental State Examination and the Hierarchic Dementia Scale.
Results	Twenty-one RCTs (N = 2253) were quantitatively analyzed. For VD, compared with Western medicine (WM), acupuncture showed better Hierarchic Dementia Scale scores ($P < 0.01$), and acupuncture plus WM also showed better Hierarchic Dementia Scale scores ($P < 0.01$). For MCI, acupuncture showed a significant improvement in Mini-Mental State Examination ($P < 0.01$) and picture recognition test scores compared with WM. For AD, WM resulted in better Hierarchic Dementia Scale scores than acupuncture ($P < 0.01$). Eight trials reported adverse events, 15 out of 2253 patients had adverse events related to acupuncture treatment, and 25 out of 2253 patients had adverse events related to WM treatment.
Conclusion	Acupuncture may be efficacious for improving cognitive function in patients with VD and MCI. However, the evidence is limited, and larger sample size and more rigorous RCTs should be conducted to verify the effectiveness and safety of acupuncture.

1.1.9. Li 2020

Li W, Wang Q, Du S, Pu Y, Xu G. Acupuncture for mild cognitive impairment in elderly people: Systematic review and meta-analyses. *Medicine (Baltimore)*. 2020;99(39). [212660]. doi

Background	Acupuncture has an unique role in preventing and managing mild cognitive impairment (MCI) in nonpharmaceutical therapies because of its small wound, mild pain, and high security for many years. However, there is no systematic review evaluating safety and efficacy of acupuncture for MCI in elderly people. Therefore, this study will provide a protocol to explore the effectiveness and safety of acupuncture for MCI in the elderly.
Methods	Retrieval from 8 electronic databases was conducted to determine eligible trials published until May, 2019. Homogeneity qualified studies were included for data were extracted such as study country location, demographic characteristics, and measure outcomes, and were analyzed by a random effect model and sensitivity analyses to identify heterogeneity. Review Manager (Revman Version 5.3) software will be used for data synthesis, sensitivity analysis, meta regression, subgroup analysis, and risk of bias assessment. A funnel plot will be developed to evaluate reporting bias.
Results	A total of 15 randomized control trials involving 1051 subjects were included . The results were as follows: Compared with the control group, the clinical efficacy rates of acupuncture was better, odds ratio = 2.52, 95% confidence interval (CI) (1.86, 3.42), $P < .00001$, mini-mental state examination scores (mean difference [MD] = 1.53, 95% CI [1.04, 2.01], $P < .00001$), Montreal cognitive assessment scores (MD = 2.05, 95% CI [1.17, 1.92], $P < .00001$), activity of daily living scale (MD = 1.71, 95% CI [-1.38, 4.79], $P > .05$), and clock drawing task scores (MD = 1.91, 95% CI [1.74, 2.08], $P < .00001$).
Conclusion	This study shows that acupuncture is beneficial for improving aspects of cognitive function in elderly people with MCI, which suggests that acupuncture may be an effective alternative and complementary approach to existing therapies for elderly people. More rigorous experimental studies and longer follow-up studies should be conducted in the future.

1.1.10. Min 2019

Min Wook Kim, Jae Hee Yoo, Hong Je Go, Seon Woo Kim, Seung Won Jang, Hyun Jin Jeong, Jae Hong Kim. Systematic Review of Acupuncture Treatment for Mild Cognitive Impairment. *Journal of Acupuncture Research*. 2019;36(2):72. [197044].

Background	The aim of this study was to review clinical efficacy of acupuncture treatment of mild cognitive impairment.
Methods	Randomized controlled trials that performed acupuncture treatment for mild cognitive impairment were retrieved from 6 online databases (PubMed, Cochrane Library, EMBASE, CNKI, NDSL, OASIS) on September 30th, 2018. Studies were selected according to inclusion and exclusion criteria, and were reviewed by Risk of Bias assessment.
Results	In total, 21 studies were included in this review. All studies were Chinese (19 studies published in Chinese and 2 in English). The sample size, 50 to 100, and the number of treatment times, 20 to 30, were the largest range in all studies. The most treatments performed was 30. The longest treatment period was 56 days, which accounted for 33% of the studies. The most frequently used evaluation index was the Mini Mental State Examination followed by the Montreal Cognitive Assessment, each used 17 times and 15 times, respectively. The most frequently used acupoints were GV20, EX-HN1, GB20, and GV24, which accounted for 47% of total number of acupoints used. In 48% of the studies, needle retention time was 30 minutes. Western medicine treatment was the most common control group. Most studies reported that the intervention group was statistically significantly different to the control group.

Conclusions	These results suggest that acupuncture for mild cognitive impairment was effective. However, it is difficult to confirm this conclusion because the quality of most of these studies were of low quality.
--------------------	---

1.1.11. Min 2016 (Vascular Cognitive Impairment) ☆

Min D, Xu-Feng W. An Updated Meta-Analysis of the Efficacy and Safety of Acupuncture Treatment for Vascular Cognitive Impairment Without Dementia. *Curr Neurovasc Res.* 2016;13(3):230-8. [164648].

Background	Vascular cognitive impairment with no dementia (VCIND) refers to a transitional state and will progress to dementia. Currently, effective medicines to control VCIND are surprisingly scarce. Acupuncture intervention is frequently advocated as an adjunct treatment for VCIND in recent studies.
Objective	So the aim of this meta-analysis was to assess the clinical efficacy and safety of acupuncture for VCIND.
Methods	The literature search was conducted in English and Chinese databases from inception until July 2015, and 15 study populations were included in the meta-analysis. Mean differences with their 95% confidence interval for continuous data were calculated with fixed effect model or random effect model using Rev Man 5.3 software.
Results	This review included 1148 subjects. The methodological qualities of the included studies were judged to be generally poor because plenty of trials had high or unclear risk of bias. The results of meta-analysis showed that acupuncture therapy was more effective in treating VCIND when compared with conventional therapy or pharmacological treatment. Two trials showed minor adverse events and two other trails mentioned no adverse events had existed.
Conclusions	Current evidences cautiously suggest that acupuncture therapy can improve the clinical efficacy for VCIND. More rigorously designed studies are needed to further confirm effectiveness and safety of acupuncture therapy in patients with VCIND.

1.1.12. Deng 2016 ☆

Deng M, Wang XF. Acupuncture for amnesic mild cognitive impairment: a meta-analysis of randomised controlled trials. *Acupunct Med.* 2016;34:342-348. [191715].

Objectives	Mild cognitive impairment (MCI) is a pre-dementia state; 5-10% of cases per year will evolve into dementia. MCI can be amnesic (AMCI) or non-amnesic. AMCI is associated with a higher risk of progression. In recent years, interest in acupuncture as a potential treatment for AMCI has grown. The aim of this meta-analysis was to estimate the clinical effectiveness and safety of acupuncture for AMCI.
Methods	Randomised controlled trials (RCTs) of acupuncture versus medical treatment for AMCI were identified using the following databases from inception to July 2015: PubMed; Medline; CENTRAL; Chinese Scientific Journal Database; The Chinese Acupuncture Trials Register; China National Knowledge Infrastructure (CNKI); and Wanfang database. Data were extracted from RCTs meeting the inclusive criteria according to Cochrane methods. Meta-analyses were conducted using Rev Man V.5.3 software.

Results	Five trials involving 568 subjects were included. Meta-analysis showed that participants receiving acupuncture had better outcomes than those receiving nimodipine with greater clinical efficacy rates (odds ratio (OR) 1.78, 95% CI 1.19 to 2.65; $p < 0.01$), mini-mental state examination (MMSE) scores (mean difference (MD) 0.99, 95% CI 0.71 to 1.28; $p < 0.01$), and picture recognition score (MD 2.12, 95% CI 1.48 to 2.75; $p < 0.01$). Meta-analysis also showed acupuncture in conjunction with nimodipine significantly improved MMSE scores (MD 1.09, 95% CI 0.29 to 1.89; $p < 0.01$) compared to nimodipine alone. Three trials reported adverse events. Methodological quality of the included studies was judged to be generally poor.
Conclusions	Acupuncture appears effective for AMCI when used as an alternative or adjunctive treatment ; however, caution must be exercised given the low methodological quality of included trials. Further, more rigorously designed studies are needed.

1.1.13. Lu 2015 (Vascular Cognitive Impairment) ☆☆

Lu Jin-Hua, Tan Jie, Zhang Hong, Yuan Yi-Qin. [Meta-analysis of acupuncture clinical efficacies on ischemic type cognitive dysfunction]. Guiding Journal of Traditional Chinese Medicine and Pharmacy. 2015;20: 87-89. [186975].

Objectives	To analysis the clinical efficacies and the adverse reactions of acupuncture treatment on ischemic type cognitive dysfunction.
Methods	The paper studied the ischemic type cognitive dysfunction patients to compare the different therapeutic effect between acupuncture and western therapy in randomized controlled trials, and to evaluate the quality of the included studies with Rev Man 5. 3 Meta-analysis software.
Results	Acupuncture treatment of ischemic type cognitive dysfunction is generally more effective than drug therapy, with summary OR=6. 12, 95%CI [4. 01, 9. 35]; In MMSE score improving respect, acupuncture treatment of ischemic type cognitive dysfunction is more effective than drug therapy, with summary MD=2. 21, 95% CI [1. 02, 3. 40]; In ADL score improving respect, acupuncture treatment is also more effective than drug therapy, with summary MD=3. 72, 95% CI [2. 80, 4. 63]. There was significant difference in Meta-analysis, and there was no significant ad-verse reactions found.
Conclusions	Acupuncture treatment of ischemic type cognitive dysfunction is more effective than drug therapy tendency, as shown in MMSE and ADL score improving respects, and the study still needs more high-quality re-searches in order to increase the strength of evidence.

1.1.14. Hu 2014 ☆☆

Hu Jun, Zhang Na, Wang Zhi-Xing, Liang Yun-Wu, Qian Qian. [Systematic review of acupuncture in treating mild cognitive impairment]. Chinese Journal of Ethnomedicine and Ethnopharmacy. 2014;12:17-21. [186940].

Objectives	Based on the principle of evidence-based medicine, the therapeutic effect and safety of acupuncture treatment for the mild cognitive impairment are systematically assessed. The Conclusion of this text would guide to the clinical practice.
-------------------	--

Methods	The 161 essays which are about acupuncture treatment for the MCI are indexed by the evidence-based medicine method. All the RCT of the essays above are estimated and 14 of them which are consistent with the internalize standard are taken into the Meta-analysis which are performed by the Review Manager 5. 0 statistical software. The sensitivity analyses and test for heterogeneity are taken at the same time. The publication bias are analysed by the funnel plot.
Results	1052 patients in 14 RCT consistent with the internalize standard are taken into the Meta-analysis, OR=2. 89, (95%CI, 2. 10~3. 97), test for overall effect Z=6. 51, P<0. 00001, indicating that there is statistical difference between the acupuncture group and the control group.
Conclusions	The acupuncture treatment for the MCI is effective and safe . However the conclusion is rather limited and it suggests that well-planned, large-scale and multi-center, double-blinded, randomized controlled trials are required.

1.1.15. Zhou 2013 (Vascular Cognitive Impairment) ☆☆

Zhou Li, Zhang Yun-Ling ,Cao Hui-Juan,And Hu Hui. [Treating vascular mild cognitive impairment by acupuncture: a systematic review of randomized controlled trials]. Chinese Journal of Integrated Traditional and Western Medicine. 2013;33(12): 1626-30. [167901].

Objectives	To systematically evaluate the effect and safety of acupuncture in the treatment of vascular mild cognitive impairment (VMCI).
Methods	Recruited were China National Knowledge Infrastructure Database (CNKI) (1979-2012), Chinese Science and Technology Periodical Database (VIP) (1989-2012), Chinese Biomedical Database (CBM), Wanfang degree and conference papers database (1985-2012) PubMed Database (1966-2012), and The Cochrane Library (issue 2012) □ The search date ended in February 2012. Randomized controlled trials (RCTs) by taking acupuncture as the main treatment for VMCI (nonvascular dementia) were collected. Results were measured using at least one internationally recognized evaluation cognitive scale. Two analysts selected the data independently. The assessment of methodological quality was based on the Cochrane Handbook and the data were analyzed by using RevMan 5. 1.0 Software. The mean difference (MD) or risk ratio (RR) were taken and graphed with 95% confidence interval (CI).
Results	Recruited 12 RCTS included a total of 691 cases meeting the inclusion criteria (all of the methodological quality was of B level). Acupuncture combined other therapies was involved in 9 RCTS, with effect compared with that of other therapies. Results of meta-analysis showed, compared with the cognitive function training alone, electroacupuncture (MD 1. 59, 95% CI 0. 69 -2. 48, P = 0. 0005, 3 studies) or body acupuncture (MD 3. 26, 95%CI 1. 69 -4. 83, P <0. 01, 1 study) combined with the cognitive function training could significantly increase the mini-mental state examination (MMSE) score of patients. In comparison to Western medicine, acupuncture could elevate ADAS-Cog score (MD 2. 16, 95%CI 1.36 -2.95, P<0.01, 3 studies). In all the studies, adverse event had not been reported.
Conclusions	Acupuncture in combination with other therapies could significantly improve cognitive functions. Acupuncture itself appeared to have better therapeutic effects than Western medicine alone.

1.1.16. Cao 2013 Ø

Cao H, Wang Y, Chang D, Zhou L, Liu J. Acupuncture for vascular mild cognitive impairment: a systematic review of randomised controlled trials. Acupuncture in Medicine. 2013;31(4):368-74. [167262].

Background	Vascular mild cognitive impairment (VMCI) is the most common type of vascular cognitive impairment induced by cerebrovascular disease. No effective medicines are currently available for VMCI.
Objective	To assess the effectiveness and safety of acupuncture for VMCI.
Methods	Seven electronic databases were searched for randomised controlled trials which investigated the effects of acupuncture compared with no treatment, placebo or conventional therapies on cognitive function or other clinical outcomes in patients with VMCI. The quality of the trials selected was evaluated according to the 'risk of bias' assessment provided by the Cochrane Handbook for Systematic Reviews of Interventions. RevMan V.5.1 software was employed for data analysis.
Results	Twelve trials with 691 participants were included. The methodological quality of all included trials was unclear and/or they had a high risk of bias. Meta-analysis showed acupuncture in conjunction with other therapies could significantly improve Mini-Mental State Examination scores (mean difference 1.99, 95% CI 1.09 to 2.88, random model, $p < 0.0001$, 6 trials). No included trials mentioned any adverse events of the treatment.
Conclusions	The current clinical evidence is not of sufficient quality for wider application of acupuncture to be recommended for the treatment of VMCI, and further large, rigorously designed trials are warranted.

1.1.17. Leung 2013 ~

Leung MCP, Yip KK, Lam CT, Lam KS, Lau W, Yu WL, Leung AKM, So KF. Acupuncture improves cognitive function (A systematic review). *Neural regeneration Research*. 2013; (18):1673-84. [165336].

Background	Acupuncture has been used as a treatment for cognitive impairment.
Objective	This review assesses clinical evidence for or against acupuncture as a treatment for cognitive impairment. This review also discusses the proposed mechanism(s) that could link acupuncture to improved cognitive function.
Methods	We searched the literature using PolyUone search from its inception to January 2013, with full text available and language limited to English. Levels of evidence were examined using Oxford Centre for Evidence-based Medicine-Levels of Evidence (March, 2009).
Results	Twelve studies met the inclusion criteria: 3 human studies and 9 animal studies. Levels of evidence ranged from level 1b to level 5.
Conclusion	Most animal studies demonstrated a positive effect of acupuncture on cognitive impairment. However, the results of human studies were inconsistent . Further high-quality human studies with greater statistical power are needed to determine the effectiveness of acupuncture and an optimal protocol.

1.2. Special Acupuncture Techniques

1.2.1. Acupuncture combined with physical therapies

1.2.1.1. Li 2021

Li RY, Huang RJ, Yu Q. Comparison of Different Physical Therapies Combined with Acupuncture for Poststroke Cognitive Impairment: a Network Meta-Analysis. *Evid Based Complement Alternat Med*. 2021 Nov 18;2021:1101101. <https://doi.org/10.1155/2021/1101101>

Objective	Physical therapy combined with acupuncture is the current research hotspot in the treatment of poststroke cognitive impairment, but which combination treatment is the best is still controversial. Based on the network meta-analysis method, we evaluated the efficacy of various physical therapies combined with acupuncture for the treatment of poststroke cognitive impairment.
Methods	We retrieved diverse randomized controlled trials of various physical therapies combined with acupuncture for the treatment of cognitive dysfunction after stroke. We selected studies, extracted data, and evaluated the risk of literature bias for the included randomized controlled trials. We used STATA 14.0 for the current network meta-analysis.
Results	Fifteen randomized controlled trials involving 1288 patients were included, which involved 7 treatment plans that included 3 control treatment plans and 4 acupuncture treatment plans combined with physical therapy. The best treatment plan for improving the Mini-Mental State Examination score of poststroke cognitive impairment is acupuncture combined with hyperbaric oxygen therapy. The best treatment option for improving the Montreal Cognitive Assessment score of poststroke cognitive impairment is acupuncture combined with hyperbaric oxygen therapy. The best option for improving the Barthel index score of poststroke cognitive impairment is acupuncture combined with transcranial magnetic stimulation. In terms of improving the overall clinical effectiveness of poststroke cognitive impairment, the best treatment option is acupuncture combined with transcranial magnetic stimulation.
Conclusion.	The analysis of all the results shows that acupuncture combined with hyperbaric oxygen therapy can significantly improve poststroke cognitive impairment compared with other combined treatments. However, due to the overall quality and quantity of the included studies, more randomized controlled trials focusing on clinical research on acupuncture combined with physical therapy for poststroke cognitive impairment are required to support the current evidence. This trial is registered with CRD42020200092.

1.2.2. Comparison of Acupuncture techniques

1.2.2.1. Yi 2024

Yi N, Xing D, Xie J, Cheng Z. Different acupuncture and moxibustion therapies for mild cognitive impairment: a network meta-analysis. Arch Gerontol Geriatr. 2024 May;120:105328.

<https://doi.org/10.1016/j.archger.2024.105328>

Objective	To investigate the effectiveness of different acupuncture and moxibustion therapies in improving cognitive function in patients with mild cognitive impairment (MCI) to determine the optimal approach. This study aims to provide insights into the treatment of MCI patients for future randomized controlled trials (RCTs) and clinical decision-making.
Materials and methods	RCT studies were retrieved from databases including PubMed, Embase, Cochrane Library, Web of Science, CNKI, WANFANG, VIP, and SinoMed. The Cochrane risk of bias tool was used to assess the risk of bias for eligible trials. Bayesian network meta-analysis was conducted using R4.2.3 software.
Results	A total of 46 RCTs with 3641 participants were included. The network meta-analysis showed that acupoint pressing + cognitive training + auricular point sticking, massage + western medicine treatment, and electroacupuncture + western medicine treatment ranked first, second, and third in improving MMSE score, respectively. Acupoint pressing + cognitive training, cognitive training + scalp acupuncture, and cognitive training + moxibustion ranked first, second, and third in improving MoCA score, respectively.

Conclusion	This study showed that acupoint pressing alone or acupoint pressing + auricular point sticking may improve cognitive function in MCI patients and possibly be the most effective acupuncture interventions for the treatment of MCI.
-------------------	--

1.2.3. Electroacupuncture

1.2.3.1. Shen 2024

Shen M, Zhang L, Li C, Ma Y, Gao S, Ma Y. Meta-analysis with trial sequential analysis investigating the impact of adjunctive electroacupuncture therapy on vascular mild cognitive impairment. *Transl Psychiatry*. 2024 Aug 30;14(1):349. <https://doi.org/10.1038/s41398-024-03052-1>

Background	To systematically collect, evaluate, and synthesize evidence from randomized controlled trials (RCTs) supporting the use of electroacupuncture (EA) as an additional treatment option for Vascular mild cognitive impairment (VaMCI), a meta-analysis was carried out.
Methods	Electronic searches of eight databases were used to locate RCTs that evaluated EA as a VaMCI adjuvant therapy. The Cochrane Risk of bias was used to assess the included trials' methodological quality. Review Manager 5.4 was used to analyze the data. Trial sequential analysis (TSA) was conducted with the trial sequential analysis program.
Results	There were 15 RCTs with 1033 subjects in them. Compared to conventional therapy (CT) alone, the Montreal Cognitive Assessment (SMD 0.72, 95 percent CI [0.55, 0.88]), Mini-mental State Examination (SMD 0.73, 95 percent CI [0.60, 0.87]), and activities of daily living (SMD 0.83, 95 percent CI [0.54, 1.12]) were significantly improved while EA was used in conjunction with CT. The current studies exceeded the required information size, according to trial sequential analysis (TSA), demonstrating the reliability of EA adjuvant therapy VaMCI.
Conclusions	According to the pooled data, EA as an adjunct therapy for the treatment of VaMCI increases clinical efficacy. Although the TSA confirms a stable conclusion, it is encouraged to conduct studies of the highest quality standards.

1.2.3.2. Kim 2019 ☆

Kim H, Kim HK, Kim SY, Kim YI, Yoo HR, Jung IC. Cognitive improvement effects of electro-acupuncture for the treatment of MCI compared with Western medications: a systematic review and Meta-analysis. *BMC Complement Altern Med*. 2019;19(1):13. [190023].

Background	Almost half of mild cognitive impairment (MCI) patients progress to dementia, which is associated with decreased quality of life and obstacles to independent living. Relevant management is expected to prevent MCI patients from progressing to dementia. In recent years, electroacupuncture (EA) has been used to treat various kinds of neurological disorders including MCI. This study evaluates the use of EA for MCI patients to increase cognitive function through a comparison with Western medications.
-------------------	--

Methods	Randomized controlled trials (RCT) or systematical reviews (SR) of EA versus Western medications for MCI were searched using the following 10 databases: Pubmed, Cochrane Library, CINAHL, EMBASE, China National Knowledge Infrastructure (CNKI), National Digital Science Library (NDSL), Journal of Oriental Neuropsychiatry (JON), Korean Medical Database (KMBASE), KoreaMed, and OASIS, from October 2007 to August 2017, without language restriction. A methodological quality assessment of RCTs or SRs that met inclusion criteria was conducted using Cochrane Risk of bias (RoB) tool and a meta-analysis by RevMan (Review Manager) 5.3.5 version of Cochrane collaboration.
Results	Five RCTs with 257 patients met inclusion criteria and those were randomly divided into two groups: the EA group (n = 103) and Western medications group (n = 154). The methodological quality of the included studies showed high risk or/and unclear of risk of bias. The meta-analysis of five studies reported that the EA group was better than the Western medications group, improving the Mini Mental State Examination (MMSE) score by 0.65 [95% CI 0.28~1.01] higher mean difference, Montreal Cognitive Assessment (MoCA) score by 0.66 [95% CI 0.00~1.32] higher mean difference. Adverse effects were not reported in the selected studies.
Conclusion	Electroacupuncture was an effective treatment for MCI patients by improving cognitive function. However, the included studies presented a low methodological quality and no adverse effects were reported. Thus, further comprehensive studies with a design in depth are needed to derive significant results.

Scalp Acupuncture

1.2.4.1. Mai 2015

Mai Wei, Zheng Jinghui. [Clinical efficacy of scalp acupuncture on mild cognitive impairment: a meta-analysis study]. Journal of Liaoning College of Traditional Chinese Medicine. 2015;12:121-124. [187007].

Objectives	To evaluate the clinical efficacy and security of scalp acupuncture on mild cognitive impairment.
Methods	Randomized Controlled trials (RCTs) concerning scalp acupuncture on mild cognitive impairment were searched from Pub Med database, VIP database, CNKI database and Wangfang database and these studies were estimated according to the Cochrane collaboration system evaluation method and were given a statistical analysis with Rev Man 5.3 software offered by the Cochrane collaboration.
Results	Five studies about MCI treated by scalp acupuncture with a total of 565 patients met inclusion criteria. The total effective rate [RR=1.53, 95%CI (1.25, 1.89), P<0.0001], effective rate [RR=14.17, 95%CI (3.44, 58.44), P=0.0002] and MMSE scores [MD=1.33, 95%CI (0.85, 1.82), P<0.00001] had a significant difference between the scalp electroacupuncture group and the nimodipine group, which showed that scalp electroacupuncture had a better effect on MCI than nimodipine. The total effective rate [RR=1.18, 95%CI (0.94, 1.47), P=0.16], effective rate [RR=1.83, 95%CI (0.72, 6.64), P=0.20] had no significant difference between the scalp acupuncture with cognitive training group and the cognitive training group, which showed that they had the same effect on MCI, while the MOCA scores [MD=2.12, 95%CI (0.78, 3.47), P=0.002] had a significant difference between the two groups which showed that scalp acupuncture with cognitive training had a better effect on MCI than cognitive training.
Conclusions	This meta-analysis suggests that scalp acupuncture has a better therapeutic efficacy on MCI and the results still need more rigorous design, large sample, multicentre randomized controlled to further verification.

1.2.5. Auricular Acupuncture

1.2.5.1. Kwon 2018

Kwon CY , Lee B , Suh HW , Chung SY , Kim JW. Efficacy and Safety of Auricular Acupuncture for Cognitive Impairment and Dementia: A Systematic Review. Evid Based Complement Alternat Med. 2018. [168729].

Objectives	To analyze the efficacy and safety of auricular acupuncture (AA) in patients with cognitive impairment and dementia.
Methods	Twelve electronic databases were searched for randomized controlled trials evaluating effects of AA in patients with cognitive impairment and/or dementia, from their inception to August 2017. The primary outcome was cognitive function, and secondary outcomes were self-care ability, quality of life, clinical efficacy rate, and incidences of adverse events.
Results	Nine studies were included, and five involving 677 participants were analyzed quantitatively. Compared with Western medications (WM), AA had mixed effects on cognitive functions (Mini-Mental State Examination [MMSE], mean difference [MD] 0.73, 95% confidence interval [CI] -0.02 to 1.48; Hierarchic Dementia Scale [HDS], MD 2.21, 95% CI 1.09 to 3.33); there was no significant improvement in the activities of daily living (ADL) score (MD 0.20, 95% CI -3.51 to 3.91) in patients with vascular dementia (VD). Compared to WM, AA combined with WM showed better clinical efficacy rate (risk ratio [RR] 1.42, 95% CI 1.06 to 1.91) in patients with VD; there was no significant improvement in cognitive functions (MMSE, MD 0.97, 95% CI -0.44 to 2.38; Montreal Cognitive Assessment [MoCA], MD 0.22, 95% CI -1.83 to 2.27) in patients with mild cognitive impairment (MCI). Compared to herbal medicine (HM), AA plus HM showed significant improvements in cognitive function (MMSE, MD 1.31, 95% CI 0.13 to 2.49) in patients with MCI and patients with vascular cognitive impairment, no dementia (VCIND) and in ADL score (MD -6.70, 95% CI -8.78 to -4.62) in patients with MCI. No adverse event associated with AA was reported.
Conclusion	The evidence reveals mixed efficacy of AA in patients with cognitive impairment and/or dementia. However, the results were inconclusive because of the small number and poor methodological quality of the included studies.

1.2.6. Acupression

1.2.6.1. Liu 2018 ☆

Liu F, Shen C, Yao L, Li Z. Acupoint Massage for Managing Cognitive Alterations in Older Adults: A Systematic Review and Meta-Analysis. Journal of Alternative and Complementary Medicine. 2018;24(6):532-540. [166427].

Background	Cognitive disorders pose a major problem in the aging population across the globe. Acupoint massage has been used to improve cognitive functions in older adults. In this study, the authors performed a meta-analysis to evaluate the usefulness of acupoint massage in preventing cognitive declines in older adults.
-------------------	---

Design	The authors searched for randomized controlled trials (RCTs) reporting on the effectiveness of acupoint massage on cognition in older adults in the following literature databases: PubMed, MEDLINE, Embase, CINAHL, ScienceDirect, Foreign Medical Journal Service, Cochrane Library, VIP Information, Chinese National Knowledge Infrastructure, WANFANG, and Chinese Biomedical. Two reviewers independently extracted the data related to the study and participants' characteristics and the cognitive impairment outcomes. Only RCTs meeting the inclusion criteria were used in the present meta-analysis.
Results	Eight RCTs with 657 participants in total (age ≥ 60 years) were included. It is actually 8 for synthetic and 6 in the meta-analysis. The authors calculated the pooled estimates of the random effects of changes in the Mini-Mental State Examination to compare the groups with and without acupoint massage. The merged mean difference (MD) was 1.94 (95% confidence interval, C.I., [1.41-2.47], $p < 0.00001$) after 3 months of acupoint massage treatment and 3.04 (95% C.I. [2.43-3.64], $p < 0.00001$) after 6 months of treatment. They also calculated the merged MD of the Wechsler Memory Scale-Revised Chinese version after 6 months of acupoint massage. Visual Reproduction was 2.95 (95% C.I. [1.30-4.60], $p = 0.0005$), Associate Learning was 1.89 (95% C.I. [1.41-2.37], $p < 0.00001$), Logical Memory was 2.85 (95% C.I. [2.06-3.63], $p < 0.00001$), and Digit Span was 3.16 (95% C.I. [2.59-3.73], $p < 0.00001$). The Cochrane Handbook for Systematic Reviews of Interventions was used to rate the quality of the studies, which was moderate overall.
Conclusion	The findings suggested that acupoint massage is an effective intervention for maintaining cognitive functions in older adults.

1.3. Special Clinical Forms

1.3.1. Poststroke Cognitive Impairment

See [corresponding item](#)

1.3.2. Sleep Disturbance in Mild Cognitive Impairment

1.3.2.1. O'Caoimh 2019

O'Caoimh R, Mannion H, Sezgin D, O'Donovan MR, Liew A, Molloy DW. Non-pharmacological treatments for sleep disturbance in mild cognitive impairment and dementia: A systematic review and meta-analysis. *Maturitas*. 2019:82-94. [200466].

Background	No disease-modifying treatments for dementia are available. Sleep disturbance is strongly associated with cognitive impairment. Non-pharmacological treatments targeting sleep may offer an alternative therapeutic approach.
Methods	We searched PubMed, CINAHL, EMBASE and the Cochrane library for non-pharmacological treatments for sleep disturbance in mild cognitive impairment (MCI) and dementia, published in English from October 1965 to 2018, including all designs, excluding studies of drug therapies.

Results	In all, 53 papers representing 48 studies were included. Participant age ranged from 67.3 to 89.4 years. Most studies (79%) had small samples (<50 participants, range 1-173) and were conducted in long-term/residential care (62%). The majority (85%) recruited participants with moderate-severe dementia; mean MMSE scores ranged from 0 to 28.3/30. Four studies examined MCI. Light therapy delivered over 1-10 weeks was the most studied stand-alone intervention (n = 27), and the majority (81.5%) of these studies found improvements on objective or subjective sleep measures, though the evidence was inconclusive with significant clinical and methodological heterogeneity. Seven multi-modal intervention studies were identified, all incorporating light exposure, and six of these reported improved sleep. Other interventions included electrotherapy stimulation (n = 4), physical exercises/activities (n = 4), acupressure/acupuncture (n = 3) and mindfulness/cognitive behavioural therapy (n = 3). Those examining MCI utilised different mono-modal approaches. A meta-analysis of data from randomised controlled trials showed a statistically significant (mean difference = 3.44, 95% CI: 0.89-5.99, I ² =0%; p = 0.008) improvement in sleep efficiency between interventions and controls, favouring the pooled interventions (bright light, multi-domain and other therapies). No other significant differences in sleep or non-sleep outcomes were found.
Conclusions	While evidence is available for non-pharmacological sleep interventions, particularly multi-domain approaches, studies were diverse and had small samples. More research examining multi-modal interventions, community-dwellers and those with MCI is required.

2. Overviews of Systematic Reviews

2.1. Han 2022

Han FH, Sun LJ, Zhang YL, Zeng ZX, Shen W, Zhan M, Wang Y, Shi JZ, Zeng XY, Lu XY, Liao X. [Overview of systematic reviews of acupuncture for vascular cognitive impairment]. Zhongguo Zhen Jiu. 2022 Jan 12;42(1):109-15. <https://doi.org/10.13703/j.0255-2930.20210207-0002>

Objective	To overview the methodological quality, report quality and evidence quality of the systematic review (SR) of acupuncture for vascular cognitive impairment (VCI).
Methods	The SRs regarding acupuncture for VCI were searched in PubMed, Cochrane Library, EMBASE, CNKI, SinoMed, Wanfang and VIP databases. The retrieval period was from the establishment of the database to September 24, 2020. The report quality, methodological quality and evidence quality of the included SRs were evaluated by PRISMA statement, the AMSTAR 2 tool and the GRADE system.
Results	A total of 22 SRs were included, including 102 outcome indexes . The methodological quality was generally low, with low scores on items 2, 5, 7, 10, 14, 15 and 16. The report quality was good, with scores ranging from 19 points to 24.5 points. The problems of report quality were mainly reflected in the aspects of structural abstract, program and registration, other analysis and funding sources. The level of outcome indexes of SRs was mostly low or very low, and the main leading factor was limitation, followed by inconsistency and inaccuracy.
Conclusion	Acupuncture for VCI is supported by low quality evidence of evidence-based medicine, but the methodological quality and evidence body quality of the relevant systematic reviews are poor, and the standardization still needs to be improved.

2.2. He 2021

He W, Li M, Han X, Zhang W. Acupuncture for Mild Cognitive Impairment and Dementia: An Overview of Systematic Reviews. *Front Aging Neurosci.* 2021. [211503]. [doi](#)

Background	Dementia is a gradual decline in cognitive ability and is becoming more common in our elderly population. Mild cognitive impairment (MCI) is defined as a slight clinical deterioration of memory capacity, below the level of normal aging, but does not constitute a clinical diagnosis of dementia. To date, no interventions have been proven to cure MCI and dementia fully. Purpose: To evaluate the potential effectiveness and safety of acupuncture for mild cognitive impairment (MCI) and dementia and evaluate the methodological quality of systematic reviews (SRs).
Methods	We conducted a literature search for SRs with meta-analyses in seven Chinese and international databases through October 1, 2020. The basic characteristics of the included SRs/meta-analyses and the basic information of the original included randomized controlled trials were extracted by three reviewers independently. A meta-analysis of the original randomized controlled trials from the included SRs/meta-analyses was performed using Stata 12.0 software. The Assessing the Methodological Quality of Systematic Reviews 2 was used to assess the methodological quality of the included SRs/meta-analyses, and the Grading of Recommendations, Assessment, Development, and Evaluation was used to rate the quality of evidence.
Results	A total of 35 SRs/meta-analyses were included, and the majority showed that acupuncture was more effective than western medicine or conventional therapy for MCI and dementia [odds ratio = 1.39; 95% confidence interval (CI): 1.24, 1.56]. There was a statistically significant difference in the Mini-Mental State Examination score (weighted mean difference = 1.23; 95% CI: 0.78, 1.68; $p < 0.00001$), and there was no significant improvement in the activities of daily living score (weighted mean difference = 1.58; 95% CI: -0.02, 3.18; $p = 0.053$). The assessment results of Assessing the Methodological Quality of Systematic Reviews 2 showed that the methodological quality of most included SRs/meta-analyses was critically low; the lowest scores were items 2, 7, and 10. For Grading of Recommendations, Assessment, Development, and Evaluation, of the 73 outcomes, 50 (68.5%) outcomes were low or very low quality, and 23 (31.5%) outcomes were moderate quality.
Conclusions	Acupuncture can be considered as an alternative for the treatment of MCI and dementia when western medicine or other therapies are contraindicated. More high-quality evidence is needed to determine further the effectiveness of acupuncture.

2.3. Hou 2020 (Methodology and Reporting Quality)

Hou T, Zheng Q, Feng X, Wang L, Liu Y, Li Y. Methodology and Reporting Quality Evaluation of Acupuncture for Mild Cognitive Impairment: An Overview of Systematic Reviews. *Evid Based Complement Alternat Med.* 2020. [211866]. [doi](#)

Objective	Since there is no consistent evidence on the effectiveness of acupuncture in the treatment of mild cognitive impairment, this review aims to summarize and critically evaluate the methodological and reporting quality of systematic reviews (SRs).
Methods	We comprehensively searched PubMed, Embase, Cochrane Library, Web of Science, China National Knowledge Infrastructure (CNKI), China Science and Technology Journal Database (VIP), Chinese Biomedical Literature (CBM), and Wanfang databases from the date of establishment to April 2019. Two authors independently selected the articles, collected the data, and assessed the identified and included SRs with the revised measurement tool to assess systematic reviews (AMSTAR 2) and preferred reporting items for SRs and meta-analyses (PRISMA). The quality of outcomes was evaluated by the Grading of Recommendations, Assessment, Development, and Evaluation (GRADE).

Results	Eleven SRs were included in this overview. The items of AMSTAR 2 in most SRs were poorly reported; only 3 SRs were rated as low quality by AMSTAR 2, and the remaining were rated as very low quality. A total of 8 SRs obtained a decent rating by PRISMA. With the GRADE tool, we have not found high-quality evidence that acupuncture is effective for mild cognitive impairment (MCI), so there is no certain conclusion on the effectiveness of acupuncture treatment for MCI.
Conclusion	The methodological and reporting quality of SRs on acupuncture for MCI is substandard, and the quality of evidence is poor. In future research, more efforts are needed to improve the quality of SRs in this field.

3. Clinical Practice Guidelines

⊕ positive recommendation (regardless of the level of evidence reported)
 ∅ negative recommendation (or lack of evidence)

3.1. Istituto Superiore di Sanità (ISS, Italy) 2024 ∅

Fabrizi E, Ancidoni A, Locuratolo N, Piscopo P, Della Gatta F, Salemme S, Pani SM, Marconi D, Vignatelli L, Sagliocca L, Caffarra P, Secreto P, Guaita A, Stracciari A, Vanacore N, Lacorte E; Guideline Working Group. The Italian guideline on diagnosis and treatment of dementia and mild cognitive impairment. *Age Ageing*. 2024 Nov 1;53(11):afae250. <https://doi.org/10.1093/ageing/afae250>

100- Do not offer acupuncture to treat cognitive symptoms in dementia. STRONG AGAINST. Adopted

From:

<https://wiki-mtc.org/> - Encyclopédie des sciences médicales chinoises

Permanent link:

<https://wiki-mtc.org/doku.php?id=acupuncture:evaluation:neuro-psychiatrie:14.%20troubles%20cognitifs>

Last update: **31 Jan 2025 15:18**